

## Stat Profile Prime Plus® Blood Gas, CO-Oximeter, Chemistry Controls Auto-Cartridge with Creatinine

Cartucho automático con creatinina para controles de química, gases en sangre y cooxímetro Stat Profile Prime Plus®, Cartouche automatique de contrôles de gaz du sang/CO-oxymètre, chimie œsphéniques avec créatine, Stat Profile Prime Plus® Auto-Kassette mit Kreatinin für Blutgas-, CO-Oximeter- und Blutchemiekontrollen, Autómatu φυσιγγίου υλικού ελέγχου χρησίας δερίνης αίματος, CO-Οξύμετρο Stat Profile Prime Plus®, Cartuccia con creatinina per controlli automatici chimici per gas ematici/CO-ossimetro Stat Profile Prime Plus®, Cartucho automático de controles de química, de CO-oxímetro e de Gás no sangue Stat Profile Prime Plus® com creatinina, Stat Profile Prime Plus® vérágz, CO-oximéter, kémiai kontrollok automatikus patron kreatininnel, Stat Profile Prime Plus® Co-Oximeter Co-oximetro 血液ガス・CO オキシメーター・生化学検査用コントロール自動カートリッジ(クレアチニン), 캐리에이티드 케라티닌 사용 Stat Profile Prime Plus® 혈액 가스, CO-산소 농도계, 화학 조절제 자동 카트리지, Stat Profile Prime Plus® 血气、一氧化碳-血氧仪、化学对照溶液自动试剂盒(含肌酐)

LOT 24043071

2025-07-23

CONTROL 1 2 3 4 5

Expected Ranges, Rangos esperados, Plages attendues, Erwartungsbereiche, Αναμενόμενα εύρηα, Intervalli previsti, Intervalos previstos, Várt tartományok, הטווחים הצפויים, 予測範囲, 예상 범위, 预期范围值

		CONTROL 1 min - $\bar{x}$ - max	CONTROL 2 min - $\bar{x}$ - max	CONTROL 3 min - $\bar{x}$ - max	CONTROL 4 min - $\bar{x}$ - max	CONTROL 5 min - $\bar{x}$ - max
pH		7.193 - 7.223 - 7.253	7.397 - 7.427 - 7.457	7.591 - 7.621 - 7.651		
H+	nmol/L	64 - 60 - 56	40 - 37 - 35	26 - 24 - 22		
PCO <sub>2</sub>	mmHg	46.6 - 53.6 - 60.6	33.6 - 38.6 - 43.6	17.3 - 21.3 - 25.3		
PCO <sub>2</sub>	kPa	6.2 - 7.1 - 8.1	4.5 - 5.1 - 5.8	2.3 - 2.8 - 3.4		
PO <sub>2</sub>	mmHg	50.6 - 60.6 - 70.6	94.3 - 104.3 - 114.3	132.6 - 147.6 - 162.6		
PO <sub>2</sub>	kPa	6.7 - 8.1 - 9.4	12.5 - 13.9 - 15.2	17.6 - 19.6 - 21.6		
SO <sub>2</sub>	%	48 - 51 - 54	77 - 80 - 83	89 - 92 - 95		
Hct	%	57 - 60 - 63	37 - 40 - 43	22 - 25 - 28		
Na <sup>+</sup>	mmol/L			137.5 - 141.5 - 145.5	111.6 - 115.6 - 119.6	
K <sup>+</sup>	mmol/L			3.69 - 3.94 - 4.19	5.92 - 6.22 - 6.52	
Cl <sup>-</sup>	mmol/L			121.3 - 125.8 - 130.3	93.3 - 97.8 - 102.3	
iCa	mmol/L			1.00 - 1.08 - 1.16	1.36 - 1.48 - 1.60	
iCa	mg/dL			4.0 - 4.3 - 4.6	5.5 - 5.9 - 6.4	
iMg	mmol/L			0.57 - 0.64 - 0.71	1.07 - 1.22 - 1.37	
iMg	mg/dL			1.4 - 1.6 - 1.7	2.6 - 3.0 - 3.3	
Glu	mg/dL			73 - 81 - 89	252 - 277 - 302	
Glu	mmol/L			4.1 - 4.5 - 4.9	14.0 - 15.4 - 16.8	
Lac	mmol/L			1.7 - 2.0 - 2.3	6.2 - 6.9 - 7.6	
Lac	mg/dL			15.1 - 17.8 - 20.5	55.2 - 61.5 - 67.7	
BUN	mg/dL			12 - 17 - 22	41 - 51 - 61	
BUN	mmol/L			4.3 - 6.1 - 7.9	14.6 - 18.2 - 21.8	
Urea	mg/dL			25.7 - 36.5 - 47.2	87.9 - 109.4 - 130.8	
Urea	mmol/L			4.3 - 6.1 - 7.9	14.6 - 18.2 - 21.8	
Creatinine	mg/dL			0.60 - 0.90 - 1.20	5.60 - 6.60 - 7.60	
Creatinine	μmol/L			0.05 - 0.08 - 0.11	0.50 - 0.58 - 0.67	
Creatinine	μmol/L			50 - 80 - 110	500 - 580 - 670	
HbF*	%	79.0 - 87.0 - 95.0	37.9 - 52.9 - 67.9	18.0 - 23.0 - 28.0		
tHb	g/dL	19.0 - 20.8 - 22.6	13.3 - 14.8 - 16.3	6.0 - 7.0 - 8.0		
tHb	g/L	190 - 208 - 226	133 - 148 - 163	60 - 70 - 80		
tHb	mmol/L	11.8 - 12.9 - 14.0	8.3 - 9.2 - 10.1	3.7 - 4.3 - 5.0		
O <sub>2</sub> Hb	%	20.0 - 22.5 - 25.0	45.3 - 49.3 - 53.3	76.0 - 81.0 - 86.0		
COHb	%	24.4 - 28.4 - 32.4	16.2 - 20.2 - 24.2	2.1 - 6.1 - 10.1		
MetHb	%	24.3 - 27.3 - 30.3	15.3 - 18.3 - 21.3	2.5 - 5.5 - 8.5		
HHb	%	17.7 - 21.7 - 25.7	8.2 - 12.2 - 16.2	3.5 - 7.5 - 11.5		
tBil*	mg/dL	17.9 - 21.9 - 25.9	9.5 - 11.5 - 13.5	5.7 - 6.1 - 6.5		
tBil*	μmol/L	306.1 - 374.5 - 442.9	162.5 - 196.7 - 230.9	97.5 - 104.3 - 111.2		
tBil*	mg/L	179.0 - 219.0 - 259.0	95.0 - 115.0 - 135.0	57.0 - 61.0 - 65.0		

EN

## Product Description

Aqueous quality control material for monitoring the performance of pH, PCO<sub>2</sub>, PO<sub>2</sub>, SO<sub>2</sub>, hematocrit (Hct), fetal hemoglobin (HbF), total hemoglobin (Hb), total bilirubin, oxygen saturation (SpO<sub>2</sub>), lactate, BUN (urea), and Creatinine in Level 4 and Levels 1, 2, and 3. A change kit consisting of 2 syringes filled with creatinine solution is included for cartridge activation prior to installation. Use of the change kit is recommended.

Intended Use

Intended for in vitro diagnostic use by healthcare professionals for monitoring the performance of the Stat Profile Prime Plus Analyzers.

Intended for in vitro diagnostic use by healthcare professionals for monitoring the performance of the Stat Profile Prime Plus Analyzers. NO RECOMMENDED USES ONLY

Storage

Store at 2-8°C (37-49°F) NOT FREEZE.

Directions for use

Discard any cartridge containing a broken or leaking syringe prior to installation of the Control Cartridge. The 2 syringes are labeled and color coded to correspond to the items on the cartridge.

Activating the cartridge

1. Hold the syringes with tip down and remove the plastic cap.
2. Attach one of the顶端的针头 to the syringe plunger and gently invert the cartridge to allow the cover to seal.
3. Remove the plastic cap from the end of the syringe and insert it into the appropriate port and insert needle.
4. Slowly depress syringe plunger until the contents are dispensed. DO NOT PULL BACK ON THE PLUNGER TO FLUSH CARTRIDGE. This will damage the plunger.
5. Remove needle/syringe assembly from barrel and discard in an appropriate sharps container.
6. Repeat Steps 1-5 for the next Control Cartridge.
7. After activating the cartridge for 1 minute, the cartridge is ready for use.

Verify that the Lot Number on the Expected Ranges Table corresponds to the Lot Number on the cartridge. Refer to Stat Profile Prime Plus Analyzer Instructions for Use Manual for complete directions.

Composition

Controls Levels 1, 2 and 3 are buffered bicarbonate solutions containing dye, salts and preservatives. Each level has a known pH and a different color. Levels 1, 2 and 3 are stable for 1 year. Controls contain dextrose, lactate, BUN (urea), creatinine and levels 1, 2 and 3 also contain Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, glucose, lactate, BUN (urea) and creatinine. Each level contains approximately 100 mL. Controls contain no constituents of human origin; however good laboratory practices should be followed during handling of these materials. (REF. NOCS DOCUMENT M29-12).

Warnings and Cautions

DO NOT FREEZE. Mix the cartridge by gently inverting for several seconds. DO NOT SHAKE CARTRIDGE. Refer to Stat Profile Prime Plus Analyzer Instructions for Use Manual for complete information.

Intended for in vitro diagnostic use. Follow standard practices for handling laboratory reagents.

Storage

Store at 2-8°C (37-49°F) NOT FREEZE.

Directions for use

Discard any cartridge containing a broken or leaking syringe prior to installation of the Control Cartridge. The 2 syringes are labeled and color coded to correspond to the items on the cartridge.

Activating the cartridge

1. Hold the syringes with tip down and remove the plastic cap.
2. Attach one of the顶端的针头 to the syringe plunger and gently invert the cartridge to allow the cover to seal.
3. Remove the plastic cap from the end of the syringe and insert it into the appropriate port and insert needle.
4. Slowly depress syringe plunger until the contents are dispensed. DO NOT PULL BACK ON THE PLUNGER TO FLUSH CARTRIDGE. This will damage the plunger.
5. Remove needle/syringe assembly from barrel and discard in an appropriate sharps container.
6. Repeat Steps 1-5 for the next Control Cartridge.
7. After activating the cartridge for 1 minute, the cartridge is ready for use.

Verify that the Lot Number on the Expected Ranges Table corresponds to the Lot Number on the cartridge. Refer to Stat Profile Prime Plus Analyzer Instructions for Use Manual for complete directions.

Limitations

PO<sub>2</sub> values are very linearly with temperature (approximately 1%°C). Therefore, it is critical to follow the temperature guidelines described in "Directions for use". The Expected range values are specific for instruments and controls manufactured by Nova Biomedical. For other instruments, refer to the Nova Profile Prime Plus Cartridge may be used for a maximum of 21 days from the initial installation date on the system at which time the system will indicate the cartridge is invalid. Each cartridge may be inserted and removed from the analyzer a maximum of 6 times.

Traceability of Standards

Total hemoglobin (Hb) and Methemoglobin (MetHb) are traceable by using Cyanmethemoglobin method. Carboxy-Hemoglobin (COHb) and Oxygen-Hemoglobin (O<sub>2</sub>Hb) are traceable using Spectrophotometry. Analyses are traced to NIST Standard Reference Materials.

Reference Intervals

Concentrations are formulated at normal and abnormal expected values in patient blood. The expected clinical range of these values in patient blood is referenced in Tietz, NW ed. 1988 Textbook of Clinical Chemistry, WB Saunders Co. Users wish to determine Mean Values and Expected Ranges in their own laboratory?

Expected Ranges

The expected range for each parameter was determined at Nova Biomedical using replicate determinations on Nova analyzers. The expected range indicates the maximum deviation from the Mean value that may be expected under differing laboratory conditions for instruments operating within specifications. Refer to Expected Ranges Table.

\*Not available for the or Point-of-Care Near-Patient Testing use.

NDC# 57841-57842-1

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and Determine Reference Intervals in the clinical laboratory, approved guideline-second edition, NCCLS C2B-A2, Volume 20, Number 13.

2How to Define and

